

Bürgi's "Kunstweg" – geometric approach

Christian Riedweg

February 29, 2016

Abstract

Placing regular $4n$ -sided polygons correctly, equations between sine and sums of sines show up – exactly these equations are used in Bürgi's method to approximate sines.

Menso Folkerts, Dieter Launert and Andreas Thom describe in [1] a method for calculating sines. This method was discovered by Jost Bürgi (1552-1632) who called it "Kunstweg". Folkerts, Launert and Thom prove that this method really works i.e. they show the convergence of this method. One of the steps in this proof is to show that for $\alpha = 90^\circ/n$ (where $n \in \mathbb{N}$)

$$\begin{pmatrix} \sin(\alpha) \\ \sin(2\alpha) \\ \sin(3\alpha) \\ \sin(4\alpha) \\ \vdots \\ \sin((n-1)\alpha) \\ 1 \end{pmatrix} \text{ is an eigenvector of } \begin{pmatrix} 1 & 1 & 1 & 1 & \dots & 1 & 1/2 \\ 1 & 2 & 2 & 2 & \dots & 2 & 2/2 \\ 1 & 2 & 3 & 3 & \dots & 3 & 3/2 \\ 1 & 2 & 3 & 4 & \dots & 4 & 4/2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 2 & 3 & 4 & \dots & n-1 & (n-1)/2 \\ 1 & 2 & 3 & 4 & \dots & n-1 & n/2 \end{pmatrix} \in \mathbb{R}^{n \times n}.$$

I.e. one has to show that for some λ the following equations hold:

$$\lambda \sin(\alpha) = \sin(\alpha) + \sin(2\alpha) + \dots + \sin((n-1)\alpha) + 1/2$$

...

$$\lambda \sin(j\alpha) = \sin(\alpha) + 2\sin(2\alpha) + \dots + j\sin(j\alpha) + j\sin((j+1)\alpha) + \dots + j\sin((n-1)\alpha) + j/2$$

...

$$\lambda = \sin(\alpha) + 2\sin(2\alpha) + 3\sin(3\alpha) + \dots + (n-1)\sin((n-1)\alpha) + n/2.$$

Subtracting subsequently the preceeding equation, this system of equations is equivalent to

$$\begin{aligned} \lambda \sin(\alpha) &= \sin(\alpha) + \sin(2\alpha) + \dots + \sin((n-1)\alpha) + 1/2 \\ \lambda \sin(2\alpha) - \lambda \sin(\alpha) &= \sin(2\alpha) + \sin(3\alpha) + \dots + \sin((n-1)\alpha) + 1/2 \\ &\dots \\ \lambda \sin(j\alpha) - \lambda \sin((j-1)\alpha) &= \sin(j\alpha) + \sin((j+1)\alpha) + \dots + \sin((n-1)\alpha) + 1/2 \\ &\dots \\ \lambda - \lambda \sin((n-1)\alpha) &= 1/2. \end{aligned} \tag{*}$$

In section 1, we develop this system of equations (*) purely geometrically (hence we give a geometric proof that the above vector is an eigenvector).

In section 2, we discuss in more details how these equations are derived from regular $4n$ -sided polygons.

For more information on Bürgi's method and further references, see [1].

1 Geometric approach to Bürgi's "Kunstweg"

Let $n \in \mathbb{N}$. We place regular convex $4n$ -sided polygons along to each other: the second not exactly on top of the first but "one step to the left". The third is placed on top of the second but "two steps to the left", and so on, as visualized below for 12-sided polygons:




Figure 1: For the $4n$ -sided polygon is $\alpha = 90^\circ / n$ (more details in the next section). Hence we need n such polygons to fill an angle of 90° at C .

The y -coordinates of the points $P_1, P_2, \dots, P_n$ are

$$R \sin(\alpha), \quad R \sin(2\alpha), \quad \dots, \quad R \sin((n-1)\alpha), \quad R.$$

Thus, the y -coordinate variation from point P_{j-1} to P_j is

$$R \sin(j\alpha) - R \sin((j-1)\alpha).$$

On the other hand, following the indicated vectors (all of length l) from P_{j-1} to P_j we get a sum of sines (more details – see next section).

Together:

$$\begin{aligned} R \sin(\alpha) &= 2l \sin(\alpha) + 2l \sin(2\alpha) + \dots + 2l \sin((n-1)\alpha) + l, \\ &\vdots \\ R \sin(j\alpha) - R \sin((j-1)\alpha) &= 2l \sin(j\alpha) + 2l \sin((j+1)\alpha) + \dots + 2l \sin((n-1)\alpha) + l \\ &\vdots \\ R - R \sin((n-1)\alpha) &= l. \end{aligned}$$

These are exactly the equations $(\star)$ that show up in Thom's proof in [1] (after dividing by $2l$ we get $\lambda = R/2l$).

2 Regular convex $4n$ -sided polygons

Let $n \in \mathbb{N}$; we consider regular convex $4n$ -sided polygons (visualized are 12-sided polygons). Let all these polygons have a side parallel to the x -axis.




Figure 2: M is the center of the regular polygon. The angle α occurs at various places. For $4n$ -sided polygons we get $\alpha = \frac{360^\circ}{4n} = \frac{90^\circ}{n}$.

Now we consider different paths along such polygons. These n paths all feature a length of **half of the polygon's circumference**, i.e. each path consists of $2n$ sides of the polygon (for regular $4n$ -sided polygons)

As we are interested in sines only, the y -coordinate of the vectors along the paths has to be considered. Thus, horizontal vectors do not contribute at all and some pairs of vectors cancel each others y -coordinate – both cases are indicated by a dashed line below.

- The **1. polygon**, first path




Figure 3: The $2n$ -vectors in the left picture are visualized in the coordinate plane (right picture). All these vectors are of the same length, say l . In Figure 2 we see that the angle between the x -axis and the first vector is α , it is 2α for the second vector, 3α for the third, and so on.

Following these vectors, the change of the y -coordinate equals

$$l \cdot \sin(\alpha) + l \cdot \sin(2\alpha) + \dots + l \cdot \sin(2n\alpha).$$

With $\alpha = \frac{90^\circ}{n}$, the considered angles are

$$\frac{90^\circ}{n}, \quad 2 \cdot \frac{90^\circ}{n}, \quad \dots, \quad n\alpha = 90^\circ, \quad \dots, \quad 2n\alpha = 180^\circ.$$

Since $\sin(90^\circ) = 1$, $\sin(180^\circ) = 0$ and $\sin(\varphi) = \sin(180^\circ - \varphi)$ we can reduce this to

$$2l \sin(\alpha) + 2l \sin(2\alpha) + \dots + 2l \sin((n-1) \cdot \alpha) + l.$$

- The j . **polygon**, where $1 < j \leq n$ (visualized is $j = 2$).




Figure 4: Let the angle of the first vector be $j\alpha$. Again, the length of the vectors is l .

Here, the angles between the x -axis and the considered vectors are

$$j\alpha, \quad (j+1)\alpha, \quad \dots, \quad (j+2n-1)\alpha.$$

Hence the y -coordinate changes by

$$l \cdot \sin(j\alpha) + l \cdot \sin((j+1)\alpha) + \dots + l \cdot \sin((j+2n-1)\alpha).$$

We can simplify this sum:

$$\begin{aligned} \text{for } j < n: & \quad 2l \sin(j\alpha) + 2l \sin((j+1)\alpha) + \dots + 2l \sin((n-1)\alpha) + l \\ \text{for } j = n: & \quad l. \end{aligned}$$

- **All polygons together:** For symmetrical reasons, the rays starting in the points P_j to the lower left (as pictured below) intersect all in one point – say C . Let C be the origin.




Figure 5: The points $P_0, P_1, \dots, P_n$ are on a circle (with center C) of some radius, say R . Note that P_n is on the y -axis since $\alpha = 90^\circ/n$.

Starting in P_0 and following the indicated path we pass the points $P_1, P_2, \dots, P_n$. By symmetry, these points are on a common circle with center C . Let R be the radius of this circle. Hence the y -coordinate of P_j is $R \sin(j\alpha)$. On the other hand, this y -coordinate is equal to the y -coordinate of P_{j-1} plus the y -coordinate change from P_{j-1} to P_j as seen above. Together we get the system of equations $(\star)$ of page 1 (after dividing by l and with $\lambda = R/l$): for P_1 :

$$R \sin(\alpha) = 2l \sin(\alpha) + 2l \sin(2\alpha) + \dots + 2l \sin((n-1)\alpha) + l,$$

for P_j , where $1 < j < n$:

$$R \sin(j\alpha) - R \sin((j-1)\alpha) = 2l \sin(j\alpha) + 2l \sin((j+1)\alpha) + \dots + 2l \sin((n-1)\alpha) + l$$

and for P_n :

$$R - R \sin((n-1)\alpha) = l.$$

References

- [1] Folkert. M. Folkerts, D. Launert, and A. Thom: *Jost Bürgi's Method for Calculating Sines*, preprint, October 2015, online: <http://arxiv.org/abs/1510.03180v2>, accessed: February 2016.